


Teacher notes

Topic E

A problem on the Compton effect

A photon of energy $m_e c^2$ (i.e. equal to the rest energy of an electron) is incident on an electron at rest. The photon scatters at an angle of 60° relative to its original direction.



Calculate

- (i) the incident photon wavelength and momentum,
- (ii) the wavelength of the scattered photon,
- (iii) the kinetic energy of the electron,
- (iv) the momentum of the electron,
- (v) the angle at which the electron moves off.

Answers

- (i) The photon wavelength and momentum are found from $\lambda = \frac{hc}{E}$; $p = \frac{h}{\lambda} \left(= \frac{E}{c} \right)$.

Hence $\lambda = \frac{hc}{E} = \frac{hc}{m_e c^2} = \frac{h}{m_e c}$. Similarly, $p = \frac{E}{c} = \frac{m_e c^2}{c} = m_e c$.

- (ii) $\lambda' - \lambda = \frac{h}{m_e c} (1 - \cos\theta) = \frac{h}{m_e c} \times \left(1 - \frac{1}{2}\right) = \frac{h}{2m_e c}$. Hence, $\lambda' = \frac{h}{2m_e c} + \frac{h}{m_e c} = \frac{3h}{2m_e c}$. (This

means that the scattered photon has energy $\frac{hc}{\lambda'} = \frac{hc}{\frac{3h}{2m_e c}} = \frac{2}{3} m_e c^2$ and momentum

$$\frac{E'}{c} = \frac{\frac{2}{3} m_e c^2}{c} = \frac{2}{3} m_e c \text{ or if you prefer } \frac{h}{\lambda'} = \frac{h}{\frac{3h}{2m_e c}} = \frac{2}{3} m_e c.)$$

- (iii) The energy transferred is $\Delta E = E - E' = m_e c^2 - \frac{2}{3} m_e c^2 = \frac{m_e c^2}{3}$.

- (iv) The mistake would be to write $\frac{m_e c^2}{3} = \frac{p^2}{2m_e}$ to get the wrong answer of $p = \sqrt{\frac{2}{3}} m_e c$.

This is wrong because the equation $E_K = \frac{p^2}{2m_e}$ is valid for non-relativistic electrons and

the electron here is relativistic. We must apply conservation of momentum to the collision of a photon with the electron.

$$m_e c = \frac{2}{3} m_e c \cos 60^\circ + p \cos \phi$$

i.e.

$$p \cos \phi = \frac{2}{3} m_e c$$

$$0 = \frac{2}{3} m_e c \sin 60^\circ - p \sin \phi$$

$$p \sin \phi = \frac{\sqrt{3}}{3} m_e c$$

This gives, squaring and adding,

$$p = \sqrt{\left(\frac{4}{9} + \frac{3}{9}\right)} m_e c = \frac{\sqrt{7}}{3} m_e c$$

(We can confirm with the relativistic formula $E^2 = (m_e c^2)^2 + p^2 c^2$. Then

$$E' = m_e c^2 + \frac{m_e c^2}{3} = \frac{4m_e c^2}{3} \text{ so that } \left(\frac{4m_e c^2}{3}\right)^2 = (m_e c^2)^2 + p^2 c^2 \text{ which solves for the}$$

momentum as $p^2 c^2 = (m_e c^2)^2 \times \left(\frac{16}{9} - 1\right) = (m_e c^2)^2 \times \frac{7}{9}$. Hence $p = \frac{\sqrt{7}}{3} m_e c$ as before.)

- (v) From

$$p \cos \varphi = \frac{2}{3} m_e c$$

$$p \sin \varphi = \frac{\sqrt{3}}{3} m_e c$$

we get, by dividing side by side,

$$\tan \varphi = \frac{\frac{\sqrt{3}}{3}}{\frac{2}{3}} = \frac{\sqrt{3}}{2} \text{ i.e. } \varphi \approx 40.9^\circ.$$